
DYSTONIA LIVING WELL SURVEY REPORT


Do you experience pain because of your dystonia?	Number of responses
Always	29%
Never	6%
Often	37%
Rarely	4%
Sometimes	25%
Grand Total	100%



9 out 10 (94%) respondents experience pain.


6 out 10 (66%) respondents are always or often in pain.

If Yes, how much pain do you experience?	Number of responses
A great deal	17%
A little	13%
A lot	31%
A moderate amount	32%
None at all	3%
(blank)	4%
Grand Total	100%




8 out of 10 (80%) respondents feel moderate to a great deal of pain.

Have you noticed sleeping worse or being more often awake at night since you have been diagnosed with dystonia (1 or more times per week)?	Number of responses
No	32%
Yes	68%
(blank)	1%
Grand Total	100%




7 out of 10 respondents (68%) don't sleep well since their dystonia diagnosis.

Do you find that you socialize less or with less pleasure because of your dystonia?	Number of responses
Always	34%
Never	6%
Rarely	8%
Sometimes	25%
Usually	27%
Grand Total	100%




6 out 10 respondents (61%) find that they usually socialize lesser.

Do you ever miss work or school because of your dystonia?	Number of responses
No	34%
Yes	55%
(blank)	11%
Grand Total	100%




More than half of the respondents (55%) miss work or school due to their dystonia.

Have you been feeling more tired or fatigued during the day since being diagnosed with dystonia?	Number of responses
Always	30%
Never	2%
Rarely	6%
Sometimes	24%
Usually	37%
(blank)	1%
Grand Total	100%




7 out of 10 respondents (68%) usually feel fatigued or tired during the day since their dystonia diagnosis.

Do you suffer from anxiety or depression since you have been diagnosed with dystonia?	Number of responses
No	28%
Yes (If yes, please specify below)	69%
(blank)	3%
Grand Total	100%



7 out of 10 respondents (69%) suffer from depression/anxiety.

Please provide details on the treatments you are currently using or have previously tried:	Number of responses
Botulinum Toxin Injections	52%
Deep Brain Stimulation (DBS)	1%
Other (please specify)	44%
(blank)	3%
Grand Total	100%



More than half of the respondents (52%) currently use Botulinum Toxin treatment

.....